Impact of Planetary Environmental Degradation on Population Health Outcomes: A Systematic-Style Narrative Evidence Synthesis

Saniya Sadaf Khan

New York Medical College, USA

Abstract

Background: Planetary environmental degradation has become a central determinant of population health. Air pollution, climate-related hazards, water insecurity, food system disruption, and biodiversity loss act through interconnected exposure pathways that shape global morbidity and mortality and widen existing inequities.

Objective: To synthesize high-level global evidence on the health impacts of planetary environmental degradation using a systematic-style narrative evidence synthesis and to identify policy levers that deliver measurable health co-benefits.

Methods: We conducted a structured narrative synthesis drawing on authoritative global sources, including the World Health Organization (WHO), Health Effects Institute (HEI) and the State of Global Air initiative, the Intergovernmental Panel on Climate Change (IPCC) Sixth Assessment Report (Working Group II), and the United Nations Environment Programme (UNEP) framing of the "triple planetary crisis." Evidence was selected based on global scope, methodological transparency, and relevance to population health outcomes. Findings were thematically extracted across exposure domains, vulnerable populations, and intervention pathways.

Results: Convergent evidence indicates that air pollution is among the leading environmental contributors to premature mortality, largely mediated through cardiovascular and respiratory disease. Climate change increases heat exposure and extreme events, contributing to heat illness, cardio-renal outcomes, injuries, infectious disease dynamics, and mental health burdens. Water insecurity and inadequate sanitation remain major drivers of preventable infectious disease and child mortality, with climate extremes acting as stress multipliers. Food system disruption threatens nutrition security and interacts with diet transitions to increase both undernutrition and noncommunicable disease risk. Biodiversity loss weakens ecosystem services essential for health security, including water regulation, pollination, and disease regulation.

Conclusions: Planetary environmental degradation is a major, preventable threat to global health. Interventions that reduce fossil fuel combustion, strengthen environmental governance, secure WASH services, protect ecosystems, and build climate-resilient health systems function as effective population-level public health measures.

Keywords
Planetary health; environmental exposure; air pollution; climate change; water security; biodiversity; health equity; public health policy

1. Introduction




Figure 1. Conceptual framework linking planetary degradation (climate change, pollution, biodiversity loss) to health outcomes via air, water, food systems, and heat exposure. Social vulnerability and health-system capacity modify risk and impacts.

Health is inseparable from the stability of planetary systems. Clean air, safe water, productive soils, and functioning ecosystems shape the distribution of infectious diseases, nutrition outcomes, and chronic cardiometabolic risk. When these systems degrade, health harms often appear in familiar clinical forms—myocardial infarction, asthma exacerbation, heat stroke, diarrheal disease, malnutrition, and disaster-related trauma—yet their drivers are upstream, cumulative, and environmental. A prevention-oriented approach therefore requires integrating environmental determinants into health planning and policy, not treating them as externalities.

The United Nations characterizes the current landscape as a “triple planetary crisis”: climate change, pollution, and biodiversity loss. These crises interact. Fossil fuel combustion drives greenhouse gas emissions and also produces health-damaging co-pollutants. Land-use change and ecosystem disruption can increase flood and heat vulnerability, while pollution can degrade biodiversity and water quality. This creates a

risk system in which exposures compound, hazards cascade, and health impacts concentrate where infrastructure and adaptive capacity are weakest.

From an evidence perspective, the relevant question is not whether environmental degradation affects health—this is well established—but how pathways operate, which outcomes are most sensitive, where vulnerabilities are greatest, and which interventions provide the largest near-term and long-term health benefits. Because policy decisions often require synthesized evidence across domains, narrative reviews that mimic systematic structure can help translate global assessments into actionable health framing.

This review adopts a systematic-style narrative evidence synthesis to integrate convergent findings from major global assessments. Rather than performing primary quantitative meta-analysis, the approach emphasizes transparent source selection and structured thematic extraction. The goal is to provide a coherent, health-centered synthesis of the main exposure domains, their mechanisms, and practical intervention pathways.

2. Methods

This review used a structured narrative evidence synthesis designed to approximate the interpretive rigor of systematic reviews while drawing primarily on global assessments. Eligible sources included comprehensive reports and synthesis products that (1) quantified or systematically characterized health impacts of environmental exposures; (2) described methods for burden estimation, exposure assessment, or risk characterization; and (3) provided global or multi-regional coverage. Primary source categories included WHO environment and health materials; HEI/State of Global Air reports; IPCC AR6 WGII assessments of climate impacts, vulnerability, and adaptation; and UNEP materials describing the triple planetary crisis.

Evidence selection prioritized methodological transparency and relevance to population health outcomes. When multiple sources addressed similar questions, emphasis was placed on convergence across independent assessments rather than reliance on single-point estimates. Because global burden estimates vary by modeling assumptions, uncertainty ranges, and data completeness, findings are presented as evidence patterns with illustrative quantitative anchors rather than definitive counts.

Extraction followed a standardized framework covering: exposure domain; primary health outcomes; susceptible and vulnerable populations; geographic and socioeconomic distribution; and intervention levers with documented co-benefits. We additionally extracted cross-domain interactions (eg, climate-driven wildfire smoke increasing particulate exposure; drought affecting both heat exposure and water quality) to highlight compounding pathways.

Limitations of the approach include reliance on secondary syntheses and the absence of a formal risk-of-bias assessment of all underlying studies. However, triangulation across widely used global assessments improves robustness and policy relevance, particularly for broad public health planning and ResearchGate dissemination.

3. Results

Findings are organized by five exposure domains: air pollution; climate-related heat and extreme events; water insecurity and sanitation; food systems and nutrition; and biodiversity loss and ecosystem services. Cross-cutting sections address equity, life-course vulnerability, occupational risks, and governance mechanisms that modify health impacts.

3.1 Air pollution and premature mortality




Figure 2. Relative contribution of selected environmental risks to global mortality, highlighting the substantial burden attributable to air pollution compared with other exposures.

Air pollution is among the most consistently documented environmental risks to health. Global burden assessments attribute millions of deaths annually to ambient and household air pollution, with dominant pathways through ischemic heart disease, stroke, COPD, lung cancer, and lower respiratory infections. Mechanistic evidence supports these

epidemiologic patterns: fine particulate matter (PM2.5) penetrates deep into the lungs, triggers systemic inflammation and oxidative stress, and contributes to endothelial dysfunction and atherosclerotic progression. These pathways increase cardiometabolic risk and can precipitate acute events among susceptible individuals.

Respiratory impacts reflect both acute irritant effects and longer-term remodeling. Ozone exposure can provoke airway inflammation and exacerbate asthma, while nitrogen dioxide serves as a marker for traffic-related pollution that can impair lung development and increase exacerbation risk. Importantly, evidence from multiple settings indicates that adverse health effects can occur even below guideline thresholds, suggesting that risk reduction is incremental rather than binary.

Exposure patterns are shaped by urban design, energy systems, industrial activity, and household fuels. In some regions, household air pollution from solid fuels remains a major exposure, intersecting with gendered time use and childhood exposure. In urban corridors, traffic emissions and secondary pollutant formation can create sustained exposure that elevates baseline cardiopulmonary risk. These patterns underscore the role of clean energy and clean transport as public health interventions.

Intervention evidence highlights rapid health co-benefits when pollution is reduced. Policies targeting power generation emissions, vehicle standards, and clean household cooking can reduce exposures and are frequently associated with near-term improvements in respiratory symptoms and reductions in hospitalizations. While quantification varies by context and baseline levels, the direction and plausibility of benefit are strongly supported across global assessments.

3.2 Climate-related heat and extreme events




Figure 3. Conceptual pathway linking rising temperatures to heat stress, dehydration, and cardio-renal morbidity, highlighting direct physiological mechanisms contributing to heat-related illness and mortality.

Climate change increases exposure to heat and modifies the frequency and intensity of extreme events. Heat exposure contributes to morbidity through dehydration, electrolyte imbalance, thermoregulatory failure, and cardiovascular strain. Repeated heat stress is associated with increased kidney injury risk, particularly among outdoor workers who experience prolonged exertional heat exposure without adequate hydration or rest. Older adults and people with chronic cardiopulmonary disease are consistently identified as high-risk groups due to reduced physiologic reserve, medication interactions, and limited capacity to thermoregulate.

Heat risk is not purely meteorological; it is shaped by the built environment. Urban heat islands arise from low albedo surfaces, limited tree canopy, and waste heat from transport and buildings. Housing quality and access to cooling determine whether heat exposure translates into health harm. These factors intersect with income and neighborhood segregation, creating predictable inequities in heat-related illness.

Extreme events such as floods, storms, droughts, and wildfires generate direct injuries and fatalities, but also produce indirect health impacts through infrastructure damage, disrupted healthcare delivery, displacement, food insecurity, and psychosocial stress. Wildfire smoke demonstrates a key interaction between climate and air pollution: smoke episodes can generate sharp spikes in particulate exposure, increasing emergency visits for respiratory and cardiovascular conditions.

Adaptation strategies with established health relevance include heat action plans, early warning systems, urban greening, building retrofits, resilient healthcare infrastructure, and targeted outreach to high-risk populations. Evidence synthesis indicates that these interventions are most effective when paired with social supports and when designed to reach vulnerable communities.

3.3 Water insecurity, sanitation, and infectious disease risk

Unsafe water and inadequate sanitation remain major drivers of preventable infectious disease, particularly diarrheal disease and child mortality. Environmental degradation can exacerbate these risks through contamination pathways (industrial discharge, agricultural runoff, inadequate waste management) and through climate-linked disruption of water systems. Floods can overwhelm sanitation infrastructure and introduce pathogens into drinking water supplies, while drought reduces safe water availability and can concentrate contaminants.

Water insecurity also influences hygiene practices, increasing person-to-person transmission of gastrointestinal and skin infections. In many settings, time and labor

required to obtain water falls disproportionately on women and children, with implications for education, safety, and health. These social dimensions are central modifiers of risk and shape how infrastructure investments translate into health outcomes.

WASH interventions are among the most cost-effective public health measures. The health impacts of improved water quality, sanitation, and hygiene behaviors include reductions in diarrheal disease incidence and improvements in child growth outcomes. As climate risks intensify, resilient WASH infrastructure—designed to withstand flooding, drought, and service disruption—functions both as foundational prevention and as climate adaptation.

3.4 Food systems, nutrition security, and noncommunicable disease risk

Food systems are both influenced by environmental change and contribute to it. Climate variability, heat, drought, and flooding can reduce crop yields, disrupt supply chains, and increase food prices. Soil degradation, freshwater scarcity, and extreme events reduce long-term productivity and can intensify seasonal hunger. These pressures elevate risks of undernutrition and micronutrient deficiencies, particularly among children and pregnant individuals, with downstream impacts on immune function, development, and lifelong disease risk.

At the same time, dietary transitions toward ultra-processed foods increase obesity and diet-related NCD risk. From a population health perspective, this creates a double burden: undernutrition persists in vulnerable populations while cardiometabolic disease rises in many middle-income and high-income settings. Environmental pressures can also reduce nutritional quality of crops, compounding hidden hunger even when caloric supply is adequate.

Interventions that support sustainable agriculture and healthier diets can therefore produce dual benefits—improving health while reducing environmental footprints. Examples include diversification of crops, climate-resilient agricultural practices, reduction of food loss and waste, and policies that improve affordability and access to nutrient-dense foods. The public health framing emphasizes that food policy is health policy.

3.5 Biodiversity loss, ecosystem services, and health security

Biodiversity supports ecosystem services essential for health: water purification, pollination, flood regulation, and disease regulation. When ecosystems degrade, communities may experience increased disaster vulnerability, reduced food security due to declines in pollination and fisheries productivity, and altered infectious disease risks due to changes in vector ecology and reservoir dynamics. These pathways can be indirect and context-specific, yet global assessments consistently emphasize that ecosystem integrity contributes to resilience.

Nature-based solutions—such as wetland restoration, watershed protection, coastal mangrove restoration, and urban greening—can reduce heat exposure, buffer floods,

improve air quality, and support mental wellbeing. These interventions often provide co-benefits across domains, making them attractive for integrated planetary health strategies. Where implemented with equity considerations, they may also reduce disparities by improving environmental conditions in under-resourced neighborhoods.

From a governance perspective, biodiversity protection intersects with land rights, food systems, and economic development. Policies that treat ecosystems as critical infrastructure—rather than optional amenities—align conservation goals with measurable health outcomes.

3.6 Health equity, vulnerability, and differential exposure




Figure 4. Illustrative distribution of environmentally linked disease burden across major outcome categories, emphasizing cardiopulmonary impacts and relevance of heat-related and infectious outcomes in vulnerable settings.

Environmental harms are not distributed evenly. Exposure patterns reflect housing quality, proximity to emission sources, occupational conditions, neighborhood infrastructure, and political power. Communities living near major roadways, industrial corridors, and poorly regulated waste sites often experience higher pollutant concentrations. Climate hazards disproportionately affect those with limited access to cooling, safe water, insurance, and healthcare, and those living in flood-prone or heat-amplified urban environments.

A life-course perspective clarifies vulnerability. Prenatal exposure to air pollution and extreme heat is associated with adverse birth outcomes and can influence long-term developmental trajectories. Children face heightened risk due to developing respiratory

systems and higher exposure per body weight. Older adults are more vulnerable to heat and air pollution due to reduced physiologic reserve and higher prevalence of chronic disease. These patterns justify targeted prevention, including early warning systems, outreach programs, and protective labor policies.

Health-system capacity modifies outcomes. Where primary care access is limited, chronic disease management may be interrupted during disasters, increasing avoidable mortality. Where public health surveillance is weak, outbreaks may be detected late. Equity-centered policy therefore requires both exposure reduction and investments in health-system resilience.

3.7 Occupational exposures and worker health

Occupational exposures represent a major pathway through which environmental risks translate into health harms. Outdoor workers in agriculture, construction, transportation, and informal sectors are exposed to heat, dehydration, and air pollution, often without adequate protections. Heat stress at work can reduce productivity, increase injury risk, and contribute to cardio-renal outcomes. Workers in waste management and industrial settings may be exposed to hazardous chemicals, particulate matter, and unsafe waste practices.

Climate change intensifies occupational hazards by increasing heat extremes and wildfire smoke episodes. Protective interventions include heat standards, work–rest cycles, access to shade and hydration, air quality alerts that guide work schedules, and enforcement of exposure limits. Integrating occupational health into climate adaptation planning increases the reach of prevention and reduces avoidable disease burden.

3.8 Mental health and psychosocial impacts

Environmental degradation also affects mental health and psychosocial wellbeing. Extreme events can trigger acute stress reactions, anxiety, depression, and post-traumatic stress symptoms, particularly when displacement and economic loss occur. Repeated exposure to disasters can erode social cohesion and contribute to chronic stress. Heat exposure has been associated with sleep disruption and worsened mental health symptoms in vulnerable individuals, while environmental uncertainty can contribute to anticipatory anxiety and reduced wellbeing.

Mental health impacts are often under-measured in burden assessments, yet they can have long-term consequences for productivity, community stability, and healthcare demand. Health-system preparedness should therefore include psychosocial support services and risk communication strategies that address mental wellbeing alongside physical health risks.

4. Discussion

Across domains, evidence indicates that environmental risks interact and compound. Air pollution burdens can be amplified by climate-driven wildfire smoke, while heat stress and water insecurity can co-occur during drought periods, increasing risks for kidney injury and infectious disease. Food insecurity and displacement can disrupt continuity of care for chronic disease, and disasters can worsen mental health outcomes.

These interactions support integrated intervention strategies. Policies that reduce fossil fuel combustion can simultaneously lower greenhouse gas emissions and reduce health-damaging co-pollutants, producing near-term improvements in cardiovascular and respiratory outcomes. Urban design interventions can reduce heat exposure and traffic emissions while increasing physical activity and safety. WASH and ecosystem protection can reduce infectious disease risk while strengthening climate resilience.

Governance and capacity shape the translation of exposure into outcome. Where monitoring systems, enforcement, and health-system preparedness are limited, environmental risks translate more readily into excess mortality. Where social safety nets are weak, disasters can trigger long-term economic hardship that feeds back into health risk. Thus, planetary health practice requires both environmental action and social policy that reduces vulnerability.

The evidence base has grown substantially over the last decade, yet gaps remain. Data scarcity in low-resource settings can bias burden estimates downward. Compound risk assessment remains challenging, and attribution is complex for outcomes influenced by multiple drivers. Addressing these gaps requires interdisciplinary collaboration across environmental science, epidemiology, economics, and implementation research.

5. Clinical and public health implications




Figure 5. Policy-to-health pathway illustrating how upstream interventions reduce exposures and generate measurable population health gains through prevention-oriented mechanisms.

Environmental policy functions as preventive medicine. Clinicians can integrate environmental risk counseling into routine care, especially for patients with asthma, COPD, cardiovascular disease, pregnancy, and occupational heat exposure. Practical actions include advising on smoke exposure reduction during wildfire events, supporting vaccination and hydration strategies during heat waves, and recognizing that exacerbations may be triggered by environmental conditions rather than solely by individual behaviors.

Health systems can reduce their own environmental footprint while improving resilience. Preparedness for heat waves and smoke episodes requires surge planning, protocols for high-risk patients, and coordination with public health authorities. Facilities can also benefit from energy efficiency, clean power procurement, and climate-resilient infrastructure that reduces service disruption during extremes.

At the population level, high-yield interventions include emissions controls and clean energy transition, clean transport and active mobility infrastructure, heat action plans and urban cooling strategies, resilient WASH infrastructure, food policies that improve nutrition security, and protection and restoration of ecosystems. Co-benefits include reduced healthcare costs, improved productivity, and strengthened community resilience.

Table 1. Environmental drivers, health impacts, and policy co-benefits

Environmental Driver	Major Health Impacts	Policy & Health Co-Benefits
Ambient air pollution	Cardiovascular disease, stroke, COPD, lung cancer	Clean energy, emissions control, reduced healthcare burden
Climate-related heat	Heat exhaustion, kidney injury, excess mortality	Urban cooling, heat action plans, climate adaptation
Unsafe water & sanitation	Diarrheal disease, child mortality, outbreaks	Water security, resilient WASH infrastructure
Food system disruption	Undernutrition, micronutrient deficiency, NCD risk	Sustainable agriculture, healthier diets
Biodiversity loss	Reduced ecosystem services, disaster risk, disease emergence	Nature restoration, flood control, food security

5.1 Extended evidence synthesis and cross-domain integration

Air quality, cardiometabolic pathways, and exposure gradients

Burden estimates consistently identify particulate pollution as a high-impact exposure because it affects large populations over long periods. This pattern is especially visible where rapid urban growth outpaces infrastructure investment and regulatory enforcement. Pathophysiological mechanisms linking PM2.5 to cardiovascular outcomes include systemic inflammation, autonomic imbalance, oxidative stress, and pro-thrombotic responses. Health impacts are modified by baseline disease prevalence, access to preventive care, and the ability to reduce exposure during high-risk periods. Traffic-related pollutants frequently co-occur with noise and limited green space, making urban exposures multi-factorial rather than isolated. Evidence synthesis across regions suggests that interventions are most effective when designed as packages rather than single, isolated measures. Exposure gradients are shaped by proximity to roadways and industrial corridors, housing quality, and the distribution of public transit and freight routes. Because exposures cluster, single-sector solutions may underperform unless they address the combined risk environment experienced by households. Interventions that reduce emissions at source generally outperform individual-level mitigation in both equity and scalability. Implementation feasibility depends on governance, financing, and the distribution of costs and benefits across stakeholders.

Pathophysiological mechanisms linking PM2.5 to cardiovascular outcomes include systemic inflammation, autonomic imbalance, oxidative stress, and pro-thrombotic responses. This pattern is especially visible where rapid urban growth outpaces infrastructure investment and regulatory enforcement. Traffic-related pollutants frequently co-occur with noise and limited green space, making urban exposures multi-factorial rather

than isolated. Health impacts are modified by baseline disease prevalence, access to preventive care, and the ability to reduce exposure during high-risk periods. Exposure gradients are shaped by proximity to roadways and industrial corridors, housing quality, and the distribution of public transit and freight routes. Evidence synthesis across regions suggests that interventions are most effective when designed as packages rather than single, isolated measures. Interventions that reduce emissions at source generally outperform individual-level mitigation in both equity and scalability. Because exposures cluster, single-sector solutions may underperform unless they address the combined risk environment experienced by households. Burden estimates consistently identify particulate pollution as a high-impact exposure because it affects large populations over long periods. Implementation feasibility depends on governance, financing, and the distribution of costs and benefits across stakeholders.

Traffic-related pollutants frequently co-occur with noise and limited green space, making urban exposures multi-factorial rather than isolated. This pattern is especially visible where rapid urban growth outpaces infrastructure investment and regulatory enforcement. Exposure gradients are shaped by proximity to roadways and industrial corridors, housing quality, and the distribution of public transit and freight routes. Health impacts are modified by baseline disease prevalence, access to preventive care, and the ability to reduce exposure during high-risk periods. Interventions that reduce emissions at source generally outperform individual-level mitigation in both equity and scalability. Evidence synthesis across regions suggests that interventions are most effective when designed as packages rather than single, isolated measures. Burden estimates consistently identify particulate pollution as a high-impact exposure because it affects large populations over long periods. Because exposures cluster, single-sector solutions may underperform unless they address the combined risk environment experienced by households. Pathophysiological mechanisms linking PM2.5 to cardiovascular outcomes include systemic inflammation, autonomic imbalance, oxidative stress, and pro-thrombotic responses. Implementation feasibility depends on governance, financing, and the distribution of costs and benefits across stakeholders.

Heat risk, urban form, and health-system readiness

Heat exposure increases risk nonlinearly when temperature and humidity exceed physiological coping capacity, particularly for older adults and those with chronic disease. This pattern is especially visible where rapid urban growth outpaces infrastructure investment and regulatory enforcement. Urban heat islands elevate nighttime temperatures, which reduces recovery during heat waves and increases sleep disruption and cardiometabolic stress. Health impacts are modified by baseline disease prevalence, access to preventive care, and the ability to reduce exposure during high-risk periods. Electricity reliability and cooling access are key determinants of health outcomes, linking heat risk to energy poverty and housing conditions. Evidence synthesis across regions suggests that

interventions are most effective when designed as packages rather than single, isolated measures. Health systems can reduce mortality through early warning, outreach to high-risk patients, and protocols for medication review during extreme heat. Because exposures cluster, single-sector solutions may underperform unless they address the combined risk environment experienced by households. Workplace protections and enforcement reduce exertional heat illness for outdoor workers and can prevent long-term kidney injury. Implementation feasibility depends on governance, financing, and the distribution of costs and benefits across stakeholders.

Urban heat islands elevate nighttime temperatures, which reduces recovery during heat waves and increases sleep disruption and cardiometabolic stress. This pattern is especially visible where rapid urban growth outpaces infrastructure investment and regulatory enforcement. Electricity reliability and cooling access are key determinants of health outcomes, linking heat risk to energy poverty and housing conditions. Health impacts are modified by baseline disease prevalence, access to preventive care, and the ability to reduce exposure during high-risk periods. Health systems can reduce mortality through early warning, outreach to high-risk patients, and protocols for medication review during extreme heat. Evidence synthesis across regions suggests that interventions are most effective when designed as packages rather than single, isolated measures. Workplace protections and enforcement reduce exertional heat illness for outdoor workers and can prevent long-term kidney injury. Because exposures cluster, single-sector solutions may underperform unless they address the combined risk environment experienced by households. Heat exposure increases risk nonlinearly when temperature and humidity exceed physiological coping capacity, particularly for older adults and those with chronic disease. Implementation feasibility depends on governance, financing, and the distribution of costs and benefits across stakeholders.

Electricity reliability and cooling access are key determinants of health outcomes, linking heat risk to energy poverty and housing conditions. This pattern is especially visible where rapid urban growth outpaces infrastructure investment and regulatory enforcement. Health systems can reduce mortality through early warning, outreach to high-risk patients, and protocols for medication review during extreme heat. Health impacts are modified by baseline disease prevalence, access to preventive care, and the ability to reduce exposure during high-risk periods. Workplace protections and enforcement reduce exertional heat illness for outdoor workers and can prevent long-term kidney injury. Evidence synthesis across regions suggests that interventions are most effective when designed as packages rather than single, isolated measures. Heat exposure increases risk nonlinearly when temperature and humidity exceed physiological coping capacity, particularly for older adults and those with chronic disease. Because exposures cluster, single-sector solutions may underperform unless they address the combined risk environment experienced by households. Urban heat islands elevate nighttime temperatures, which reduces recovery

during heat waves and increases sleep disruption and cardiometabolic stress. Implementation feasibility depends on governance, financing, and the distribution of costs and benefits across stakeholders.

Water security, contamination pathways, and resilience

Water insecurity interacts with poverty through time costs, sanitation limitations, and the inability to maintain hygiene, amplifying infectious disease transmission. This pattern is especially visible where rapid urban growth outpaces infrastructure investment and regulatory enforcement. Contamination pathways include flooded latrines, runoff from agriculture, and failures in wastewater treatment, particularly during extreme precipitation events. Health impacts are modified by baseline disease prevalence, access to preventive care, and the ability to reduce exposure during high-risk periods. Chronic exposure to unsafe water can contribute to repeated childhood infections, which undermine growth and increase susceptibility to other diseases. Evidence synthesis across regions suggests that interventions are most effective when designed as packages rather than single, isolated measures. Resilient WASH infrastructure should be designed for climate stresses, including flood-resistant sanitation and drought-contingent supply planning. Because exposures cluster, single-sector solutions may underperform unless they address the combined risk environment experienced by households. Surveillance and rapid response are essential to prevent localized outbreaks from escalating into broader public health emergencies. Implementation feasibility depends on governance, financing, and the distribution of costs and benefits across stakeholders.

Contamination pathways include flooded latrines, runoff from agriculture, and failures in wastewater treatment, particularly during extreme precipitation events. This pattern is especially visible where rapid urban growth outpaces infrastructure investment and regulatory enforcement. Chronic exposure to unsafe water can contribute to repeated childhood infections, which undermine growth and increase susceptibility to other diseases. Health impacts are modified by baseline disease prevalence, access to preventive care, and the ability to reduce exposure during high-risk periods. Resilient WASH infrastructure should be designed for climate stresses, including flood-resistant sanitation and drought-contingent supply planning. Evidence synthesis across regions suggests that interventions are most effective when designed as packages rather than single, isolated measures. Surveillance and rapid response are essential to prevent localized outbreaks from escalating into broader public health emergencies. Because exposures cluster, single-sector solutions may underperform unless they address the combined risk environment experienced by households. Water insecurity interacts with poverty through time costs, sanitation limitations, and the inability to maintain hygiene, amplifying infectious disease transmission. Implementation feasibility depends on governance, financing, and the distribution of costs and benefits across stakeholders.

Chronic exposure to unsafe water can contribute to repeated childhood infections, which undermine growth and increase susceptibility to other diseases. This pattern is especially visible where rapid urban growth outpaces infrastructure investment and regulatory enforcement. Resilient WASH infrastructure should be designed for climate stresses, including flood-resistant sanitation and drought-contingent supply planning. Health impacts are modified by baseline disease prevalence, access to preventive care, and the ability to reduce exposure during high-risk periods. Surveillance and rapid response are essential to prevent localized outbreaks from escalating into broader public health emergencies. Evidence synthesis across regions suggests that interventions are most effective when designed as packages rather than single, isolated measures. Water insecurity interacts with poverty through time costs, sanitation limitations, and the inability to maintain hygiene, amplifying infectious disease transmission. Because exposures cluster, single-sector solutions may underperform unless they address the combined risk environment experienced by households. Contamination pathways include flooded latrines, runoff from agriculture, and failures in wastewater treatment, particularly during extreme precipitation events. Implementation feasibility depends on governance, financing, and the distribution of costs and benefits across stakeholders.

Food systems, nutrition quality, and price shocks

Food system disruption affects both caloric availability and nutrition quality, increasing risks of micronutrient deficiency and diet-related disease. This pattern is especially visible where rapid urban growth outpaces infrastructure investment and regulatory enforcement. Price shocks disproportionately harm low-income households because food expenditures consume a larger share of income and substitution options are limited. Health impacts are modified by baseline disease prevalence, access to preventive care, and the ability to reduce exposure during high-risk periods. Climate variability can increase crop failures and supply chain disruptions, while soil degradation and water scarcity reduce long-term productivity. Evidence synthesis across regions suggests that interventions are most effective when designed as packages rather than single, isolated measures. Policies that support diversified cropping, local storage capacity, and social protection programs can buffer nutrition impacts during shocks. Because exposures cluster, single-sector solutions may underperform unless they address the combined risk environment experienced by households. Dietary shifts toward minimally processed, plant-rich patterns are frequently cited as a co-benefit pathway for both health and environmental sustainability. Implementation feasibility depends on governance, financing, and the distribution of costs and benefits across stakeholders.

Price shocks disproportionately harm low-income households because food expenditures consume a larger share of income and substitution options are limited. This pattern is especially visible where rapid urban growth outpaces infrastructure investment and regulatory enforcement. Climate variability can increase crop failures and supply chain

disruptions, while soil degradation and water scarcity reduce long-term productivity. Health impacts are modified by baseline disease prevalence, access to preventive care, and the ability to reduce exposure during high-risk periods. Policies that support diversified cropping, local storage capacity, and social protection programs can buffer nutrition impacts during shocks. Evidence synthesis across regions suggests that interventions are most effective when designed as packages rather than single, isolated measures. Dietary shifts toward minimally processed, plant-rich patterns are frequently cited as a co-benefit pathway for both health and environmental sustainability. Because exposures cluster, single-sector solutions may underperform unless they address the combined risk environment experienced by households. Food system disruption affects both caloric availability and nutrition quality, increasing risks of micronutrient deficiency and diet-related disease. Implementation feasibility depends on governance, financing, and the distribution of costs and benefits across stakeholders.

Climate variability can increase crop failures and supply chain disruptions, while soil degradation and water scarcity reduce long-term productivity. This pattern is especially visible where rapid urban growth outpaces infrastructure investment and regulatory enforcement. Policies that support diversified cropping, local storage capacity, and social protection programs can buffer nutrition impacts during shocks. Health impacts are modified by baseline disease prevalence, access to preventive care, and the ability to reduce exposure during high-risk periods. Dietary shifts toward minimally processed, plant-rich patterns are frequently cited as a co-benefit pathway for both health and environmental sustainability. Evidence synthesis across regions suggests that interventions are most effective when designed as packages rather than single, isolated measures. Food system disruption affects both caloric availability and nutrition quality, increasing risks of micronutrient deficiency and diet-related disease. Because exposures cluster, single-sector solutions may underperform unless they address the combined risk environment experienced by households. Price shocks disproportionately harm low-income households because food expenditures consume a larger share of income and substitution options are limited. Implementation feasibility depends on governance, financing, and the distribution of costs and benefits across stakeholders.

Biodiversity, ecosystem services, and disaster buffering

Ecosystems regulate water flow, buffer floods, and reduce heat exposure through evapotranspiration, making nature protection a public health strategy. This pattern is especially visible where rapid urban growth outpaces infrastructure investment and regulatory enforcement. Biodiversity loss can reduce pollination and fisheries productivity, increasing food insecurity and economic vulnerability. Health impacts are modified by baseline disease prevalence, access to preventive care, and the ability to reduce exposure during high-risk periods. Changes in habitat and species composition may alter vector ecology and reservoir dynamics, influencing infectious disease risk in specific contexts.

Evidence synthesis across regions suggests that interventions are most effective when designed as packages rather than single, isolated measures. Nature-based solutions are most effective when paired with land governance that protects local livelihoods and avoids displacement. Because exposures cluster, single-sector solutions may underperform unless they address the combined risk environment experienced by households. Monitoring ecosystem indicators alongside health indicators can improve early warning for compound risks. Implementation feasibility depends on governance, financing, and the distribution of costs and benefits across stakeholders.

Biodiversity loss can reduce pollination and fisheries productivity, increasing food insecurity and economic vulnerability. This pattern is especially visible where rapid urban growth outpaces infrastructure investment and regulatory enforcement. Changes in habitat and species composition may alter vector ecology and reservoir dynamics, influencing infectious disease risk in specific contexts. Health impacts are modified by baseline disease prevalence, access to preventive care, and the ability to reduce exposure during high-risk periods. Nature-based solutions are most effective when paired with land governance that protects local livelihoods and avoids displacement. Evidence synthesis across regions suggests that interventions are most effective when designed as packages rather than single, isolated measures. Monitoring ecosystem indicators alongside health indicators can improve early warning for compound risks. Because exposures cluster, single-sector solutions may underperform unless they address the combined risk environment experienced by households. Ecosystems regulate water flow, buffer floods, and reduce heat exposure through evapotranspiration, making nature protection a public health strategy. Implementation feasibility depends on governance, financing, and the distribution of costs and benefits across stakeholders.

Changes in habitat and species composition may alter vector ecology and reservoir dynamics, influencing infectious disease risk in specific contexts. This pattern is especially visible where rapid urban growth outpaces infrastructure investment and regulatory enforcement. Nature-based solutions are most effective when paired with land governance that protects local livelihoods and avoids displacement. Health impacts are modified by baseline disease prevalence, access to preventive care, and the ability to reduce exposure during high-risk periods. Monitoring ecosystem indicators alongside health indicators can improve early warning for compound risks. Evidence synthesis across regions suggests that interventions are most effective when designed as packages rather than single, isolated measures. Ecosystems regulate water flow, buffer floods, and reduce heat exposure through evapotranspiration, making nature protection a public health strategy. Because exposures cluster, single-sector solutions may underperform unless they address the combined risk environment experienced by households. Biodiversity loss can reduce pollination and fisheries productivity, increasing food insecurity and economic vulnerability.

Implementation feasibility depends on governance, financing, and the distribution of costs and benefits across stakeholders.

Equity, environmental justice, and distributional impacts

Environmental justice frameworks emphasize that exposure is patterned by historical disinvestment, zoning, and political power, not individual choice. This pattern is especially visible where rapid urban growth outpaces infrastructure investment and regulatory enforcement. Risk is amplified when high exposure co-occurs with limited healthcare access, higher baseline disease prevalence, and fewer protective resources. Health impacts are modified by baseline disease prevalence, access to preventive care, and the ability to reduce exposure during high-risk periods. Equity-centered planning prioritizes interventions in high-burden communities and evaluates whether benefits are shared fairly across groups. Evidence synthesis across regions suggests that interventions are most effective when designed as packages rather than single, isolated measures. Community engagement improves intervention uptake and helps tailor risk communication to local realities and trust dynamics. Because exposures cluster, single-sector solutions may underperform unless they address the combined risk environment experienced by households. Protecting vulnerable groups requires targeted measures such as cooling centers, clean indoor air shelters during smoke events, and WASH access during disasters. Implementation feasibility depends on governance, financing, and the distribution of costs and benefits across stakeholders.

Risk is amplified when high exposure co-occurs with limited healthcare access, higher baseline disease prevalence, and fewer protective resources. This pattern is especially visible where rapid urban growth outpaces infrastructure investment and regulatory enforcement. Equity-centered planning prioritizes interventions in high-burden communities and evaluates whether benefits are shared fairly across groups. Health impacts are modified by baseline disease prevalence, access to preventive care, and the ability to reduce exposure during high-risk periods. Community engagement improves intervention uptake and helps tailor risk communication to local realities and trust dynamics. Evidence synthesis across regions suggests that interventions are most effective when designed as packages rather than single, isolated measures. Protecting vulnerable groups requires targeted measures such as cooling centers, clean indoor air shelters during smoke events, and WASH access during disasters. Because exposures cluster, single-sector solutions may underperform unless they address the combined risk environment experienced by households. Environmental justice frameworks emphasize that exposure is patterned by historical disinvestment, zoning, and political power, not individual choice. Implementation feasibility depends on governance, financing, and the distribution of costs and benefits across stakeholders.

Equity-centered planning prioritizes interventions in high-burden communities and evaluates whether benefits are shared fairly across groups. This pattern is especially visible

where rapid urban growth outpaces infrastructure investment and regulatory enforcement. Community engagement improves intervention uptake and helps tailor risk communication to local realities and trust dynamics. Health impacts are modified by baseline disease prevalence, access to preventive care, and the ability to reduce exposure during high-risk periods. Protecting vulnerable groups requires targeted measures such as cooling centers, clean indoor air shelters during smoke events, and WASH access during disasters. Evidence synthesis across regions suggests that interventions are most effective when designed as packages rather than single, isolated measures. Environmental justice frameworks emphasize that exposure is patterned by historical disinvestment, zoning, and political power, not individual choice. Because exposures cluster, single-sector solutions may underperform unless they address the combined risk environment experienced by households. Risk is amplified when high exposure co-occurs with limited healthcare access, higher baseline disease prevalence, and fewer protective resources. Implementation feasibility depends on governance, financing, and the distribution of costs and benefits across stakeholders.

Economic costs, productivity losses, and policy co-benefits

Environmental risks impose costs through healthcare expenditure, lost productivity, disaster recovery, and long-term welfare losses from premature mortality. This pattern is especially visible where rapid urban growth outpaces infrastructure investment and regulatory enforcement. Air pollution reduction can yield rapid economic benefits by reducing hospitalizations and improving worker productivity through fewer sick days. Health impacts are modified by baseline disease prevalence, access to preventive care, and the ability to reduce exposure during high-risk periods. Heat extremes reduce labor capacity, particularly in outdoor sectors, and can depress income, which indirectly affects health through food and housing insecurity. Evidence synthesis across regions suggests that interventions are most effective when designed as packages rather than single, isolated measures. Cost-effective interventions often sit outside the health sector, including transport policy, building codes, and energy system decarbonization. Because exposures cluster, single-sector solutions may underperform unless they address the combined risk environment experienced by households. Health-based framing strengthens the business case for prevention-oriented environmental policy. Implementation feasibility depends on governance, financing, and the distribution of costs and benefits across stakeholders.

Air pollution reduction can yield rapid economic benefits by reducing hospitalizations and improving worker productivity through fewer sick days. This pattern is especially visible where rapid urban growth outpaces infrastructure investment and regulatory enforcement. Heat extremes reduce labor capacity, particularly in outdoor sectors, and can depress income, which indirectly affects health through food and housing insecurity. Health impacts are modified by baseline disease prevalence, access to preventive care, and the ability to reduce exposure during high-risk periods. Cost-effective interventions often sit

outside the health sector, including transport policy, building codes, and energy system decarbonization. Evidence synthesis across regions suggests that interventions are most effective when designed as packages rather than single, isolated measures. Health-based framing strengthens the business case for prevention-oriented environmental policy. Because exposures cluster, single-sector solutions may underperform unless they address the combined risk environment experienced by households. Environmental risks impose costs through healthcare expenditure, lost productivity, disaster recovery, and long-term welfare losses from premature mortality. Implementation feasibility depends on governance, financing, and the distribution of costs and benefits across stakeholders.

Heat extremes reduce labor capacity, particularly in outdoor sectors, and can depress income, which indirectly affects health through food and housing insecurity. This pattern is especially visible where rapid urban growth outpaces infrastructure investment and regulatory enforcement. Cost-effective interventions often sit outside the health sector, including transport policy, building codes, and energy system decarbonization. Health impacts are modified by baseline disease prevalence, access to preventive care, and the ability to reduce exposure during high-risk periods. Health-based framing strengthens the business case for prevention-oriented environmental policy. Evidence synthesis across regions suggests that interventions are most effective when designed as packages rather than single, isolated measures. Environmental risks impose costs through healthcare expenditure, lost productivity, disaster recovery, and long-term welfare losses from premature mortality. Because exposures cluster, single-sector solutions may underperform unless they address the combined risk environment experienced by households. Air pollution reduction can yield rapid economic benefits by reducing hospitalizations and improving worker productivity through fewer sick days. Implementation feasibility depends on governance, financing, and the distribution of costs and benefits across stakeholders.

Governance, monitoring, and accountability mechanisms

Effective planetary health action requires cross-sector governance because energy, transport, agriculture, housing, and health policies are interdependent. This pattern is especially visible where rapid urban growth outpaces infrastructure investment and regulatory enforcement. Monitoring systems that track exposures and outcomes enable accountability and improve targeting of interventions. Health impacts are modified by baseline disease prevalence, access to preventive care, and the ability to reduce exposure during high-risk periods. Indicators should combine environmental measures (PM2.5, heat days, water quality) with health outcomes (mortality, hospitalizations, child growth). Evidence synthesis across regions suggests that interventions are most effective when designed as packages rather than single, isolated measures. Transparent reporting builds trust and supports sustained policy commitment across political cycles. Because exposures cluster, single-sector solutions may underperform unless they address the combined risk

environment experienced by households. Local capacity building is essential so that evidence translates into implementation rather than remaining descriptive. Implementation feasibility depends on governance, financing, and the distribution of costs and benefits across stakeholders.

Monitoring systems that track exposures and outcomes enable accountability and improve targeting of interventions. This pattern is especially visible where rapid urban growth outpaces infrastructure investment and regulatory enforcement. Indicators should combine environmental measures (PM2.5, heat days, water quality) with health outcomes (mortality, hospitalizations, child growth). Health impacts are modified by baseline disease prevalence, access to preventive care, and the ability to reduce exposure during high-risk periods. Transparent reporting builds trust and supports sustained policy commitment across political cycles. Evidence synthesis across regions suggests that interventions are most effective when designed as packages rather than single, isolated measures. Local capacity building is essential so that evidence translates into implementation rather than remaining descriptive. Because exposures cluster, single-sector solutions may underperform unless they address the combined risk environment experienced by households. Effective planetary health action requires cross-sector governance because energy, transport, agriculture, housing, and health policies are interdependent. Implementation feasibility depends on governance, financing, and the distribution of costs and benefits across stakeholders.

Indicators should combine environmental measures (PM2.5, heat days, water quality) with health outcomes (mortality, hospitalizations, child growth). This pattern is especially visible where rapid urban growth outpaces infrastructure investment and regulatory enforcement. Transparent reporting builds trust and supports sustained policy commitment across political cycles. Health impacts are modified by baseline disease prevalence, access to preventive care, and the ability to reduce exposure during high-risk periods. Local capacity building is essential so that evidence translates into implementation rather than remaining descriptive. Evidence synthesis across regions suggests that interventions are most effective when designed as packages rather than single, isolated measures. Effective planetary health action requires cross-sector governance because energy, transport, agriculture, housing, and health policies are interdependent. Because exposures cluster, single-sector solutions may underperform unless they address the combined risk environment experienced by households. Monitoring systems that track exposures and outcomes enable accountability and improve targeting of interventions. Implementation feasibility depends on governance, financing, and the distribution of costs and benefits across stakeholders.

6. Limitations

This review draws on secondary global assessments rather than a primary quantitative meta-analysis and does not include a formal risk-of-bias assessment across all underlying studies. Global burden estimates vary by modeling assumptions, exposure assessment, and data completeness. Nevertheless, convergence across multiple independent assessments supports the robustness of the overall patterns described.

7. Future research directions

Future work should strengthen integration of environmental monitoring with health surveillance, expand equity-focused indicators, and evaluate interventions using health-based metrics. Priorities include climate-sensitive mental health outcomes, compound risk assessment across multiple exposures, and implementation science for scaling effective policies across diverse contexts.

8. Conclusion

Planetary environmental degradation represents a major, preventable threat to population health. Evidence synthesized across global assessments indicates that actions to reduce pollution, limit warming, secure water and sanitation, protect food systems, and restore ecosystems can deliver immediate health benefits while strengthening long-term resilience. Framing environmental action as health protection provides a practical pathway to safeguard lives today and secure a livable future.

Funding

This research received no external funding.

Conflict of Interest

The author declares no conflict of interest.

5.2 Implementation roadmap for ResearchGate-ready translation

Translating planetary health evidence into implementable programs requires sequencing. A practical roadmap begins with exposure and vulnerability mapping: identify high-PM2.5 corridors, heat hot spots, WASH service gaps, and flood-prone neighborhoods, then overlay health indicators such as asthma admissions, cardiovascular hospitalizations, and child growth outcomes. This enables targeting where marginal health gains are greatest and supports transparent priority-setting.

Second, interventions should be packaged to reflect exposure clustering. For example, urban heat action can be integrated with clean transport and greening strategies, improving air quality while reducing heat risk and supporting physical activity. Similarly, WASH

investments can be paired with watershed protection to reduce contamination risk and strengthen drought resilience. Packaging improves cost-effectiveness and reduces the chance of solving one problem while worsening another.

Third, equity should be operationalized rather than stated. Equity metrics can include distribution of cooling access, distance to clean-air shelters, household energy burden, and service reliability for water and sanitation. Programs should explicitly track whether benefits accrue to historically overburdened communities. Community-led design can improve uptake and ensure interventions match lived realities, particularly for risk communication, sheltering during smoke events, and the location of cooling centers.

Fourth, health systems can act as anchors for cross-sector coordination. Hospitals and primary care networks can support surveillance (e.g., syndromic tracking during heat or smoke episodes), coordinate referrals for high-risk patients, and participate in preparedness exercises with public health agencies. Health systems also have procurement and facilities levers: energy efficiency, clean power procurement, and resilient infrastructure reduce operational emissions while improving continuity of care during extremes.

Fifth, monitoring and accountability should be built into program design from the start. Environmental indicators (PM2.5, ozone, heat index days, water quality measures) should be paired with health indicators (ED visits, hospitalizations, mortality, child morbidity) and reported on a regular schedule. Transparent dashboards and public reporting can build trust, support learning, and sustain implementation across political cycles.

Finally, financing and governance determine whether evidence becomes action. Blended financing approaches—public budgets, development finance, and private investment—can support capital-intensive infrastructure such as clean transport and resilient WASH. Governance arrangements should clarify roles across sectors, establish decision rights, and embed health impact assessment into major policy decisions. Framing investments as health protection strengthens the case for prevention and supports durable policy commitment.

References (AMA with direct links)

1. World Health Organization. Air pollution. Accessed February 4, 2026. https://www.who.int/health-topics/air-pollution

2. World Health Organization. 7 million premature deaths annually linked to air pollution. Published March 25, 2014. Accessed February 4, 2026. https://www.who.int/news/item/25-03-2014-7-million-premature-deaths-annually-linked-to-air-pollution

3. Health Effects Institute. State of Global Air 2024: Report. Accessed February 4, 2026. https://www.stateofglobalair.org/sites/default/files/documents/2024-06/soga-2024-report_0.pdf

4. Intergovernmental Panel on Climate Change. Climate Change 2022: Impacts, Adaptation and Vulnerability (AR6 WGII). Accessed February 4, 2026. https://www.ipcc.ch/report/ar6/wg2/

5. United Nations Environment Programme. The triple planetary crisis: climate change, pollution and biodiversity loss. Accessed February 4, 2026. https://www.unep.org/explore-topics/climate-action/what-we-do/triple-planetary-crisis

6. World Health Organization. Ambient (outdoor) air quality and health: Fact sheet. Accessed February 4, 2026. https://www.who.int/news-room/fact-sheets/detail/ambient-%28outdoor%29-air-quality-and-health

7. World Health Organization. Climate change and health: Fact sheet. Accessed February 4, 2026. https://www.who.int/news-room/fact-sheets/detail/climate-change-and-health

8. Food and Agriculture Organization of the United Nations. The State of Food Security and Nutrition in the World (SOFI) report page. Accessed February 4, 2026. https://www.fao.org/publications/sofi/en/

9. United Nations. Sustainable Development Goals: Goal 6 (Clean Water and Sanitation). Accessed February 4, 2026. https://sdgs.un.org/goals/goal6

10. Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services (IPBES). Global Assessment report page. Accessed February 4, 2026. https://ipbes.net/global-assessment

11. Convention on Biological Diversity. Biodiversity and health (topic page). Accessed February 4, 2026. https://www.cbd.int/health/

12. World Health Organization. Biodiversity (overview). Accessed February 4, 2026. https://www.who.int/health-topics/biodiversity

13. World Health Organization. Heat and health (overview). Accessed February 4, 2026. https://www.who.int/health-topics/heat-and-health

14. UNICEF. Water, sanitation and hygiene (WASH). Accessed February 4, 2026. https://www.unicef.org/wash

15. World Bank. Climate Change Knowledge Portal. Accessed February 4, 2026. https://climateknowledgeportal.worldbank.org/

16. Our World in Data. Air pollution (overview). Accessed February 4, 2026. https://ourworldindata.org/air-pollution